

Selective C-H bond electro-oxidation of benzylic acetates and alcohols to benzaldehydes

Barone, Mateus R; Jones, Alan M

DOI:
[10.1039/C7OB02300F](https://doi.org/10.1039/C7OB02300F)

License:
Creative Commons: Attribution (CC BY)

Document Version
Peer reviewed version

Citation for published version (Harvard):
Barone, MR & Jones, AM 2017, 'Selective C-H bond electro-oxidation of benzylic acetates and alcohols to benzaldehydes', *Organic and Biomolecular Chemistry*, vol. 15, no. 47, pp. 10010-10015.
<https://doi.org/10.1039/C7OB02300F>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:
DOI: 10.1039/C7OB02300F. Citation: Org. Biomol. Chem., 2017,15, 10010-10015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.


If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: M. R. Barone and A. M. Jones, *Org. Biomol. Chem.*, 2017, DOI: 10.1039/C7OB02300F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Selective C-H Bond Electro-oxidation of Benzylic Acetates and Alcohols to Benzaldehydes

Mateus R. Barone^{ab} and Alan M. Jones^{c*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A chemical oxidant-free and mediator-free, direct electro-oxidation of both benzylic alcohols and benzylic esters are reported. The scope of the reaction is explored as a function of both steric and electronic effects. Expansion of the scope to non-benzylic and heteroaryl substrates are investigated. Functionalisation of esters and alcohols selectively to the aldehyde oxidation level using a traceless electron approach is reported.

Introduction

The controlled and chemo-selective oxidation of primary alcohols to aldehydes and secondary alcohols to the corresponding ketones are fundamental reactions in organic synthesis.¹ In turn, these aldehydes and ketones serve as precursors for a variety of complexity generating reactions.² However, the controlled oxidation of primary alcohols to aldehydes can be problematic due to over-oxidation to the carboxylic acid oxidation state.³ There are a range of versatile chemical oxidants available to the academic and industrial chemist that enable this transformation to be performed on demand.⁴ However, the majority of these oxidation reactions are stoichiometric in nature and therefore suffer from the generation of quantities of chemical waste.⁵

Recently, the field of electrosynthesis has undergone a renaissance⁶ and has found application in a variety of organic synthetic transformations, such as: C-H bond activation,⁷ total synthesis,⁸ Diels-Alder reaction⁹ amongst others.¹⁰ A fundamental advantage of the electrosynthesis approach is the replacement of the need to use stoichiometric oxidants and instead the oxidation reaction is performed on the electrode surface *via* quantum mechanical tunnelling¹¹ or through a mediator in solution.¹²

To address the challenge of identifying a cleaner oxidation, we explored the use of electrosynthesis to replace the need for both a chemical oxidant and mediator in these REDOX transformations. The simultaneous removal of oxidant and

mediator would minimise chemical waste associated with the reaction. We have recently investigated the Shono-type oxidation of C-H bonds adjacent to a tertiary amide¹³ and our initial foray into this area began with attempting to expand the scope of the amide oxidation to esters (Figure 1). It is known that the Shono electro-oxidation of amides proceeds through an *N*-acyl iminium species (**A**).¹⁴ It was therefore postulated that a transient *O*-acyloxonium species (**B**)¹⁵ could form under similar electro-oxidative conditions in esters bearing an α -methylene or methide group.




Figure 1. Context: Expanding the scope of amide C-H bond functionalisation from amides (**A**) to esters (**B**).

This unstable *O*-acyloxonium species (**B**) would in turn react further with adventitious water to form the aldehyde product and a carboxylic acid as a by-product *via* intermediate (**C**). Alternatively, an ester and aldehyde product could potentially form when conducted in an alcohol. To the best of our knowledge there are only limited reports of electro-oxidative cleavage of an ester group.¹⁶

^a Department of Chemical Engineering, University of Sao Paulo, Brazil

^b Division of Chemistry and Environmental Science, Manchester Metropolitan University, M1 5GD, UK

^c School of Pharmacy, University of Birmingham, Edgbaston, B15 2TT, UK

* (A.M.J.) E-mail: a.m.jones.2@bham.ac.uk

Electronic Supplementary Information (ESI) available: copies of ¹H and ¹³C NMR spectra and cyclic voltammetry measurements (PDF). See DOI: 10.1039/x0xx00000x



ARTICLE

Journal Name

Results and discussion

To probe whether this methodological leap was indeed feasible, a collection of benzylic acetates were prepared including the parent benzyl acetate **1a**, a mild electron donating example **1b** and a mild electron withdrawing example **1c** (structures shown in Table 2). Cyclic voltammetry was recorded for **1a** (Figure 2).




Figure 2. Cyclic voltammetry measurement on substrate **1a** (5 mM) at a scan rate of 10 mVs⁻¹ in 9:1 acetonitrile:methanol and TBAP (0.5 M) using an RVC working electrode, RVC counter electrode and a Ag/AgCl reference electrode..

Using a sensitive measurement of 10 mVs⁻¹ scan rate it was observed in both electrolyte systems (LiClO₄ and Bu₄NClO₄)¹⁷ that oxidation waves for all three substrates were observed with +1.0 V (**1a**), +1.2 V (**1b**) and +0.9 V (**1c**) oxidation potentials (relative to Ag/AgCl) and a slightly improved peak current (*i*_p) measurement in Bu₄NClO₄. On the basis of this positive oxidation result, screening of potential conditions to enable viable electro-oxidation of model benzylic acetate **1a** was attempted (Table 1).

In the first instance, potentiostatic conditions were screened with **1a** (entries 1-8) using both Bu₄NClO₄ and LiClO₄ as the electrolyte. The applied voltage was varied around the observed oxidation potential for **1a** (cyclic voltammetry measurement, +1.0 V) with up to an extra 300 mV applied to compensate for expected iR drop across the electrode surface.¹⁸ In all cases, the reaction was performed until Fmol⁻¹ equivalent to 4 electrons per mole of substrate was passed or




Table 2. Electro-oxidation of benzylic and homologated esters (n.r. = no reaction, c.e. = current efficiency).

starting material consumption was observed. Near the oxidation potential of **1a** trace conversion to the desired aldehyde **2a** was observed in both electrolyte systems (entries 3-4 and 6-7, respectively). However, at higher applied voltages degradation products were observed (entries 5 and 8,

Entry	Electrolyte	Solvent	Voltage (mV)	Current (mA)	Conversion (%)
1	Bu ₄ NClO ₄	MeCN/MeOH	+900	-	0
2	Bu ₄ NClO ₄	MeCN/MeOH	+1000	-	0
3	Bu ₄ NClO ₄	MeCN/MeOH	+1100	-	<5
4	Bu ₄ NClO ₄	MeCN/MeOH	+1200	-	<5
5	Bu ₄ NClO ₄	MeCN/MeOH	+1300	-	degradation
6	LiClO ₄	MeCN/MeOH	+1100	-	<10
7	LiClO ₄	MeCN/MeOH	+1200	-	<10
8	LiClO ₄	MeCN/MeOH	+1300	-	degradation
9	Bu ₄ NClO ₄	MeCN/MeOH	-	20	27
10	Bu ₄ NClO ₄	MeOH	-	20	34
11	Bu ₄ NClO ₄	MeOH	-	10	49
12	LiClO ₄	MeCN/MeOH	-	20	25
13	LiClO ₄	MeOH	-	20	57
14	LiClO ₄	MeOH	-	10	84
15	LiClO ₄	MeOH/H ₂ O	-	10	38
16	LiClO ₄	MeOH	-	-	0

Table 1. Reaction condition optimisation for **1a** to **2a**.



Journal Name

ARTICLE

approach was respectively), coupled with excessive time required for sufficient charge to be passed (24 h to 5 days), a controlled voltage ruled out in this system early on. Switching to a galvanostatic approach, using the same electrolyte systems and reticulated vitreous carbon (RVC) electrodes produced more promising results (entries 9-15).

In particular, it was observed that a single solvent system gave higher yields than the previously optimized solvent system for the Shono oxidation,¹³ acetonitrile-methanol (entries 9 vs 10 and 12 vs 13, respectively). Lowering the current density across the electrode surface from an applied 20 mA to 10 mA, led to a doubling in reaction time (approx. 5-

6 h) and a concomitant improvement in reaction yield in both electrolyte systems (entries 11 vs 10 and 14 vs 13, respectively). Lowering the current rate further led to modest improvements in conversion but unacceptable lengthening of the reaction time. It became clear that lithium perchlorate was superior to tetrabutylammonium perchlorate (entry 14 vs 11) plus coupled with its ease of separation from organic products was selected as the electrolyte. To probe whether additional water improved the reaction yield (entry 15) based on a postulated mechanism led to a reduced yield versus entry 14. Furthermore, passing no electrical current led to no reaction (entry 16). To test the hypothesis that C-H bond




Table 3. Results of direct alcohol electro-oxidation to aldehydes (c.e. = current efficiency).

oxidation adjacent to an ester was possible, a collection of benzylic esters were prepared. Our initial results using the optimised procedure are detailed in Table 2. It was found using the optimised conditions, appreciable amounts of the desired aldehyde (**2a-2c**) were obtained from the benzylic acetates (**1a-1c**). To address, the mechanism issue identified in Figure 1,


a simple aqueous base wash removed the acid by-product. However, there were still limitations to this approach for example electro-oxidation of cyclic benzylic ester **1d** or homologated ester **1e**, did not afford the aldehyde **2d** and **2e**, respectively. These limitations coupled with the use of an acyl ancillary group still did not meet our green chemistry



standards due to the additional manipulation step required to prepare the acetate. We therefore considered whether a stabilising group on the heteroatom adjacent to the C-H bond was indeed essential to electro-oxidation. Promisingly, cyclic voltammetry measurements on benzyl alcohol **3a** showed an oxidation wave at +1.2 V relative to Ag/AgCl.¹⁹ Using our previously optimised conditions for the ester electro-oxidation we explored the direct mediator-free oxidation on a range of commercially available alcohols (Table 3).

It was possible to cleanly convert benzyl alcohol (**3a**) to benzaldehyde (**2a**) under the mild conditions of lithium perchlorate in methanol in near quantitative yield and in improved yield compared to the chemical manipulation of forming the benzyl acetate (99% vs 80%, respectively).²⁰ The scope of this reaction was further investigated *via* exploration of the effects of electron withdrawing and donating group around the ring system and steric effects around the reacting centre. The use of a chlorine atom as an electron withdrawing group in the *ortho*- (**2f**, 81%), *meta*- (**2g**, 55%) and *para*- (**2h**, 86%) positions was well tolerated. The use of a methyl group as a mild electron donating group in the *ortho*- (**2i**, 66%), *meta*- (**2j**, 99%) and *para*- (**2b**, 83%) positions also afforded good to excellent yields of the aldehyde.

Intriguingly, a strong electron donating group *para* to the reacting centre (from *para*-methoxy benzyl alcohol, **3i** (not shown)) resulted in a greatly reduced yield of aldehyde **2k** (27%). The relatively low yield of **2k** compared with other benzaldehydes is likely to be due to the *para*-methoxy group stabilizing the *carbon*-centred benzylic radical leading to further unproductive reaction pathways and oxidative decomposition of **2k**.



Scheme 1. Proposed mechanisms for (a) benzyl alcohol and (b) benzyl acetate electro-oxidation (w/u = work-up).

The result for a strongly electron withdrawing group in the 4-position of the substrate **3j** afforded only trace conversion to the desired product as a result of the highly electron withdrawing nature of the *para*-nitro group. Based on the experimental results above and EPR literature²¹ of related systems a tentative mechanism can be proposed (Scheme 1).

Electro-oxidation of the benzyl alcohol reveals a benzyl radical, subsequent oxidation of the hydroxyl group will deliver an aldehyde after deprotonation. Similarly, the benzylic acetate system could proceed via analogous initial steps to an *O*-acyloxonium species that could be trapped with water or methanol to deliver a hemiacetal or acetal species, respectively. In turn, the hemi-acetal or acetal intermediate would collapse upon work-up to deliver the aldehyde (Scheme 1B). We have shown that mild electron withdrawing and donating groups are tolerated in both systems but those that strongly donate or withdraw electrons were less compatible with direct electro-oxidation. Nitrogen containing heterocycle (**3n**) and non-benzylic substrates (**3k-3m**) gave trace conversion (<5%) to the aldehyde. However, changing the benzene ring to a thiophene was successful, affording aldehyde **2h** in quantitative yield, demonstrating the utility of the approach in other classes of heterocyclic systems.

Conclusion

The use of mediator-free, galvanostatic, direct electro-oxidation of benzylic alcohols and acetates shows scope on a range of substrates and offers an alternative and complementary approach to the preparation of valuable aldehydes. Mechanistic investigations to determine the sequence of electro-oxidation steps are now underway.

Experimental section

General methods

Reactions were carried out under nitrogen. Organic solutions were dried over MgSO₄. Starting materials were purchased from commercial suppliers and were used without further purification. Solvents were dried over molecular sieves (3-4Å). Flash silica chromatography was performed using Sigma-Aldrich high-purity grade, pore size 60 Å, 200-400 mesh particle size silica gel. ¹H and ¹³C NMR spectra were recorded on a JEOL ECS 400 NMR Spectrometer at 400 MHz or Bruker AVIII 300 or 400 MHz spectrometers. Chemical shifts (δ) are reported relative to TMS (δ=0) and/or referenced to the solvent in which they were measured. All chemical shifts (δ) are reported in parts per million, and coupling constants (J) are reported in Hertz. Low and High-resolution mass spectrometry analysis were obtained using an Agilent 6450 LC-MS/MS system. Infrared (IR) spectra were recorded on a ThermoScientific Nicolet Impact-380 ATR-FTIR spectrometer. Electrosynthesis and cyclic voltammetry were performed using a Metrohm Autolab PGSTAT100N + 10 Amp Booster (Metrohm Autolab, U.K.) with an undivided electrochemical cell. Data was processed using Autolab Nova software (version 2.0). The electrode system for electrosynthesis consisted of two reticulated vitreous carbon (RVC) electrodes (supplier: ERG Aerospace; specifications: 45 pores per inch (PPI); 800 square foot per cubic foot; size 20 x 11 x 5 mm) for galvanostatic reactions plus an additional *pseudo* Ag/AgCl reference



electrode for potentiostatic reactions. Please see¹³ for further details and images of the reaction set-up.

General procedures and safety statements

(A) Preparation of Benzyl acetates²²

A round bottom flask equipped with a magnetic stirrer was charged with the appropriate benzyl alcohol (10.0 mmol) and acetic anhydride (15.0 mmol). The reaction mixture was refluxed (85 °C) for 15 h. Upon cooling to room temperature the reaction mixture was poured into water (50 mL). The aqueous phase was extracted with ethyl acetate (30 mL x 3). The combined organic phase was washed with 1 M aq HCl (30 mL), saturated aq NaHCO₃ (30 mL), water (30 mL) and brine (30 mL), and dried (MgSO₄). The solvent was evaporated under reduced pressure to afford the title compound without further purification.

(B) Electrosynthetic oxidation reactions

A sealable electrochemical cell equipped with a magnetic stirrer was added anhydrous methanol (10 mL), Lithium perchlorate (0.53 g [0.50 M]) and the alcohol (0.5 mmol [0.05 M]) or the benzyl acetate (0.05 M) under study. Two reticulated vitreous carbon (RVC) electrodes were inserted into the solution at a distance of approximately 0.5 cm from each other and the vial sealed. The solution was cooled to 0 °C (ice-bath) and degassed with nitrogen. A fixed current (10 mA) was passed through the solution in an ice bath until the desired charge (*Q*) was achieved (on average *t* = 5.5 h). Upon completion of the reaction, the solvent was evaporated (at 35 °C), the residual oil was partitioned between water (10 mL) and ethyl acetate (5 mL x 3). The combined organic extracts were dried (MgSO₄) and evaporated. Analytical quality sample of the aldehyde/ketone was obtained by column chromatography (SiO₂; cyclohexane-ethyl acetate) to afford the title compound.

(C) Cyclic voltammetry studies

An undivided glass cell equipped with a rectangular reticulated vitreous carbon (RVC) anode (11 cm²) and rectangular RVC cathode (11 cm²), arranged opposite to one another at a distance of 3.0 mm with a Ag/AgCl *pseudo* reference electrode placed 1.0 mm from the working electrode. To this reaction vessel was added the analyte under study (5.0 mM) in the organic solvent(s) (total volume 6.0 mL) and electrolyte (e.g. tetrabutylammonium perchlorate (0.5 M)). Scan rate was varied using the Autolab Nova 2.0 software.

CAUTION: Perchlorates – May intensify fire. In combination with flammable solvents, all sources of ignition should be avoided. During the concentration stage of the work-up, low temperature rotary evaporation with a suitable shield should be employed. The aqueous layer (after extraction) was treated with excess sodium sulphite prior to disposal.

CAUTION: Passing an electrical current through flammable organic solvent – The potentiostat and wiring should be regularly inspected for faults and pass all relevant electrical tests prior to use.

Spectroscopic data

Benzyl acetate (1a)²³

The title compound was afforded as a clear oil (0.85 g, 56%) using general procedure A. ¹H NMR (400 MHz, CDCl₃) δ 7.42–7.29 (m, 5H), 5.12 (s, 2H), 2.11 (s, 3H) ppm; ¹³C NMR (100.1 MHz, CDCl₃) δ 171.0, 136.0, 128.7, 128.4, 128.3, 66.4, 21.1 ppm; *m/z* (ESI) = 151 [M+H]⁺. Spectroscopic data are in accordance with literature.

4-Methylbenzyl acetate (1b)²⁴

The title compound was afforded as a clear oil (0.78 g, 48%) using general procedure A. ¹H NMR (400 MHz, CDCl₃) δ 7.34–7.23 (m, 2H), 7.23–7.10 (m, 2H), 5.07 (s, 2H), 2.36 (s, 3H), 2.08 (s, 3H) ppm; ¹³C NMR (100.1 MHz, CDCl₃) δ 171.1, 138.2, 133.0, 129.3, 128.5, 66.4, 21.3, 21.2 ppm; *m/z* (ESI) = 165 [M+H]⁺. Spectroscopic data are in accordance with literature.

4-Chlorobenzyl acetate (1c)²⁵

The title compound was afforded as a clear oil (0.98 g, 53%) using general procedure A. ¹H NMR (400 MHz, CDCl₃) δ 7.34–7.30 (m, 2H), 7.30–7.25 (m, 2H), 5.05 (s, 2H), 2.08 (s, 3H) ppm; ¹³C NMR (100.1 MHz, CDCl₃) δ 170.9, 134.5, 134.2, 129.7, 128.8, 65.6, 21.1 ppm; *m/z* (ESI) = 185 [M³⁵Cl+H]⁺, 187 [M³⁷Cl+H]⁺. Spectroscopic data are in accordance with literature.

Phenylethyl acetate (1e)²⁶

The title compound was afforded as a clear oil (0.58 g, 36%) using general procedure A. ¹H NMR (400 MHz, CDCl₃) δ 7.34–7.19 (m, 5H), 4.27 (t, *J* = 7.0 Hz, 2H), 2.99–2.83 (t, *J* = 7.0 Hz, 2H), 2.03 (s, 3H) ppm; *m/z* (ESI) = 165 [M+H]⁺. Spectroscopic data are in accordance with literature.

Benzaldehyde (2a)²⁷

The title compound was afforded as a colourless oil (52 mg, 99%) using general procedure B (from benzyl alcohol, **3a**). Alternatively, the title compound (42 mg, 80%) was prepared using general procedure B (from benzyl acetate, **1a**). ¹H NMR (300 MHz, CD₃OD) δ 10.01 (s, 1H), 7.91–7.85 (m, 2H), 7.68–7.63 (m, 1H), 7.60–7.45 (m, 2H) ppm; ¹³C NMR (100.1 MHz, CD₃OD) δ 192.6, 136.4, 134.6, 129.9, 129.1 ppm; *m/z* (ESI) 106 [M]⁺. Spectroscopic data are in accordance with literature.

4-Methylbenzaldehyde (2b)²⁸

The title compound was afforded as a colourless oil (50 mg, 83%) using general procedure B (from 4-methylbenzyl alcohol, **3b**). Alternatively, the title compound (90 mg, 75%) was prepared using general procedure B (from 4-methylbenzyl acetate, **1b**). ¹H NMR (300 MHz, CD₃OD) δ 9.95 (s, 1H), 7.77 (d, *J* = 8.0 Hz, 2H), 7.32 (d, *J* = 8.0 Hz, 2H), 2.43 (s, 3H) ppm; ¹³C NMR (100.1 MHz, CD₃OD) δ 191.9, 145.5, 134.2, 129.8, 129.7, 21.7 ppm; *m/z* (ESI) 121 [M+H]⁺. Spectroscopic data are in accordance with literature.

4-Chlorobenzaldehyde (2c)²⁸

The title compound was afforded as a white powder (60 mg, 86%) using general procedure B (from 4-chlorobenzyl alcohol, **3c**). Alternatively, the title compound (48 mg, 68%) was prepared using general procedure A (from 4-chlorobenzyl acetate, **1c**). ¹H NMR (300 MHz, CDCl₃) δ 10.00 (s, 1H), 7.94–7.78 (m, 2H), 7.71–7.43 (m, 2H) ppm; ¹³C NMR (100.1 MHz, CDCl₃) δ 190.1, 140.7, 134.5, 130.1, 129.2 ppm; *m/z* (ESI) 139 [M-H]⁺. Spectroscopic data are in accordance with literature.

2-Chlorobenzaldehyde (2f)²⁸



ARTICLE

Journal Name

The title compound was afforded as a colourless oil (57 mg, 81%) using general procedure B (from 2-chlorobenzyl alcohol, **3d**). ^1H NMR (300 MHz, CDCl_3) δ 10.48 (s, 1H), 7.92-7.89 (m, 1H), 7.56-7.50 (m, 1H), 7.49-7.42 (m, 2H) ppm; ^{13}C NMR (100.1 MHz, CDCl_3) δ ppm 190.0, 138.2, 135.2, 132.5, 130.7, 129.5, 127.4 ppm; m/z (ESI) 139 $[\text{M}-\text{H}]^+$; Hi-Res LC-MS (ESI) m/z calcd for $\text{C}_7\text{H}_5\text{OCl}$ $[\text{M}-\text{H}]^+$ 138.9956, found 138.9961. Spectroscopic data are in accordance with literature.

3-Chlorobenzaldehyde (2g)²⁸

The title compound was afforded as a colourless oil (39 mg, 55%) using general procedure B (from 3-chlorobenzyl alcohol). ^1H NMR (300 MHz, CDCl_3) δ 9.97 (s, 1H) 7.89-7.83 (m, 1H) 7.79-7.73 (m, 1H) 7.51-7.45 (m, 2H) ppm; ^{13}C NMR (100.1 MHz, CDCl_3) δ 190.8, 137.8, 135.4, 134.5, 130.5, 129.4, 127.8 ppm; m/z (ESI) 139 $[\text{M}-\text{H}]^+$. Spectroscopic data are in accordance with literature.

Thiophene-2-carbaldehyde (2h)²⁹

The title compound was afforded as a pale yellow oil (55 mg, 99%) using general procedure B (from 2-thiophenemethanol). ^1H NMR (400 MHz, CDCl_3) δ 9.95 (d, J = 1.3 Hz, 1H), 7.78 (ddd, J = 5.0, 3.8, 1.2 Hz, 2H), 7.22 (dd, J = 5.0, 3.8 Hz, 1H) ppm; ^{13}C NMR (100.1 MHz, CDCl_3) δ 183.0, 144.0, 136.3, 135.1, 128.3 ppm; m/z (ESI) 111 $[\text{M}-\text{H}]^+$. Spectroscopic data are in accordance with literature.

2-Methylbenzaldehyde (2i)²⁷

The title compound was afforded as a colourless oil (39 mg, 65%) using general procedure B (from 2-methylbenzyl alcohol). ^1H NMR (300 MHz, CDCl_3) δ 10.29 (s, 1H), 7.82 (d, J = 7.3 Hz, 1H), 7.51-7.31 (m, 3H), 2.69 (s, 3H) ppm; ^{13}C NMR (100.1 MHz, CDCl_3) δ 192.7, 140.2, 133.9, 133.5, 131.9, 131.5, 126.1, 20.1 ppm; m/z (ESI) 121 $[\text{M}+\text{H}]^+$; Hi-Res LC-MS (ESI) m/z calcd for $\text{C}_8\text{H}_8\text{O}$ $[\text{M}+\text{H}]^+$ 121.0648, found 121.0644. Spectroscopic data are in accordance with literature.

3-Methylbenzaldehyde (2j)³⁰

The title compound was afforded as a clear oil (59 mg, 99%) using general procedure B (from 3-methylbenzyl alcohol). ^1H NMR (300 MHz, CDCl_3) δ 9.98 (s, 1H), 7.77-7.58 (m, 2H) 7.48-7.38 (m, 2H), 2.43 (s, 3H) ppm; ^{13}C NMR (100.1 MHz, CDCl_3) δ 192.8, 138.4, 136.5, 135.4, 130.1, 129.0, 127.3, 21.5 ppm; m/z (ESI) 121 $[\text{M}+\text{H}]^+$. Spectroscopic data are in accordance with literature.

4-Methoxybenzaldehyde (2k)²⁷

The title compound was afforded as a colourless oil (18 mg, 27%) using general procedure B (from 4-methoxybenzyl alcohol). ^1H NMR (300 MHz, CDCl_3) δ ppm 9.88 (s, 1H) 7.88-7.78 (m, 2H) 7.09-6.95 (m, 2H) 3.89 (s, 3H); ^{13}C NMR (100.1 MHz, CDCl_3) δ 190.4, 164.3, 131.5, 129.8, 113.9, 55.6 ppm; m/z (ESI) 135 $[\text{M}-\text{H}]^+$; Hi-Res LC-MS (ESI) m/z calcd for $\text{C}_8\text{H}_8\text{O}_2$ $[\text{M}-\text{H}]^+$; 135.0452, found 135.0454. Spectroscopic data are in accordance with literature.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

A.M.J. thanks the Royal Society of London (United Kingdom) for the award of a research grant (RG150135). The authors thank the analytical services at Manchester Metropolitan University and the Centre for Chemical and Materials Analysis in the School of Chemistry at the University of Birmingham for analytical support. The NMR instruments used in this research were obtained through Birmingham Science City: Innovative Uses for Advanced Materials in the Modern World (West Midlands Centre for Advanced Materials Project 2), with support from Advantage West Midlands (AWM) and part funded by the European Regional Development Fund (ERDF).

Notes and references

- (a) G. Tojo, M. I. Fernandez, *Oxidation of Alcohols to Aldehydes and Ketones: A Guide to Current Common Practice*, Springer, New York, 2006; (b) G. Franz, R. A. Sheldon, *Oxidation*, in: *Ullmann's Encyclopaedia of Industrial Chemistry*, VCH-Verlag, Weinheim, 2000; (c) R. C. Larock, *Comprehensive Organic Transformations*, 2nd Ed, Wiley, New York, 1999.
- (a) D. Lee, J. K. Sello, S. L. Schreiber, *Org. Lett.*, 2000, **2**, 709; (b) A. F. Abdel-Magid, S. J. Mehrman, *Org. Process Res. Dev.*, 2006, **10**, 971.
- M. Sankar, E. Nowicka, E. Carter, D. M. Murphy, D. W. Knight, D. Bethell, G. J. Hutchings, *Nature Comm.*, 2014, **5**, 3332.
- (a) D. Könnig, T. Olbrisch, F. D. Sypaseuth, C. C. Tzschucke, M. Christmann, *Chem. Commun.*, 2014, **50**, 5014; (b) J. E. Steves, S. S. Stahl, *J. Am. Chem. Soc.*, 2013, **135**, 15742; (c) S. B. Ley, J. Norman, W. P. Griffith, S. P. Marsden, *Synthesis*, 1994, 639.
- A. N. Campbell, S. S. Stahl, *Acc. Chem. Res.*, 2012, **45**, 851.
- (a) E. J. Horn, B. R. Rosen, P. S. Baran, *ACS Cent. Sci.*, 2016, **2**, 302; (b) E. J. Horn, B. R. Rosen, Y. Chen, J. Tang, K. Chen, M. D. Eastgate, P. S. Baran, *Nature*, 2016, **533**, 78; (c) M. Yan, Y. Kawamata, P. S. Baran, *Chem. Rev.* doi: 10.1021/acs.chemrev.7b00397
- (a) S. Liang, C.-C. Zeng, H.-Y. Tian, B.-G. Sun, X.-G. Luo, F. Z. Ren, *J. Org. Chem.*, 2016, **81**, 11565; (b) R. Hayashi, A. Shimizu, Y. Song, Y. Ashikari, T. Nokami, J.-I. Yoshida, *Chem. Eur. J.*, 2016, **23**, 61; (c) Y. Kawamata, M. Yan, Z. Liu, D.-H. Bao, J. Chen, J. Starr, P. S. Baran, *J. Am. Chem. Soc.*, 2017, **139**, 7448; (d) A. G. O'Brien, A. Maruyama, Y. Inokuma, M. Fujita, P. S. Baran, *Angew. Chem. Int. Ed.*, 2014, **53**, 11868.
- B. R. Rosen, E. W. Werner, A. G. O'Brien, P. S. Baran, *J. Am. Chem. Soc.*, 2014, **136**, 5571.
- J. C. Moore, E. S. Davies, D. A. Walsh, P. Sharmaab, J. E. Moses, *Chem. Commun.*, 2014, **50**, 12523.
- P. Du, J. L. Brosmer, D. G. Peters, *Org. Lett.*, 2011, **13**, 4072.
- S. G. Christov, *Ber. Bunsenges. Phys. Chem.*, 1975, **79**, 357.
- (a) R. Ciriminna, M. Ghahremani, B. Karimi, M. Pagliaro, *Chemistry Open*, 2017, **6**, 5; (b) A. Badalyan, S. S. Stahl, *Nature*, 2016, **535**, 406; (c) C.-B. Bai, N.-X. Wang, X.-W. Lan, Y.-J. Wang, Y. Xing, J.-L. Wen, X.-W. Gao, W. Zhang, *Sci. Rep.*, 2016, **6**, 20163.
- (a) A. M. Jones, C. E. Banks, *Beilstein J. Org. Chem.*, 2014, **10**, 3056; (b) P. Alfonso-Suárez, A. V. Kolliopoulos, J. P. Smith, C. E. Banks, A. M. Jones, *Tetrahedron Lett.*, 2015, **56**, 6863.
- T. Shono, Y. Matsumura, K. Tsubata, *J. Am. Chem. Soc.*, 1981, **103**, 1172.
- A. K. Ghosh, S. S. Kulkarni, C.-X. Xu, K. Shurrush, *Tetrahedron Asymm.*, 2008, **19**, 1020.



- 16 (a) E. Baciocchi, M. Mattioli, R. Romano, R. Ruzziconi, *J. Org. Chem.*, 1991, **56**, 7154; (b) C.-C. Zeng, J. Y. Becker, *J. Org. Chem.* 2004, **69**, 1053; (c) E. A. Mayeda, L. L. Miller, J. F. Wolf, *J. Am. Chem. Soc.* 1972, **94**, 6812; (d) H. Lund, *Acta Chem. Scand.* 1957, **11**, 491.
- 17 Scan rate sweeps for **1a-1c** can be found in the supporting information.
- 18 W. Oelßner, F. Berthold, U. Guth, *Mater. Corros.*, 2006, **57**, 455.
- 19 Cyclic voltammetry measurements of **3a** are in the accompanying supporting information.
- 20 Exposure of **3a** to lithium perchlorate in methanol without passing an electric current did not afford any **2a** after 12 h.
- 21 L. Meng, J. Su, Z. Zha, L. Zhang, Z. Zhang, Z. Wang, *Chem. Eur. J.* 2013, **19**, 5542.
- 22 J. M. Álvarez-Calero, Z. D. Jorge, G. M. Massanet, *Org. Lett.*, 2016, **18**, 6344.
- 23 B. Janza, A. Studer, *J. Org. Chem.*, 2005, **70**, 6991.
- 24 T. Mino, Y. Hasegawa, M. Sakamoto, T. Fujita, *J. Organomet. Chem.* 2007, **692**, 4389.
- 25 J. W. J. Bosco, A. K. Saikia, *Chem. Commun.*, **2004**, 1116.
- 26 C. B. Rao, D. C. Rao, D. C., Babu, Y. Venkateswarlu, *Eur. J. Org. Chem.*, 2010, **15**, 2855.
- 27 C. K. Lin, T.-J. Lu, *Tetrahedron*, 2010, **66**, 9688-9693.
- 28 K. Moriyama, M. Takemura, H. Togo, *J. Org. Chem.*, 2014, **79**, 6094.
- 29 S. Imai, H. Togo, *Tetrahedron*, 2016, **72**, 6948.
- 30 A. Kamimura, H. Komatsu, T. Moriyama, Y. Nozaki, *Tetrahedron*, 2013, **69**, 5968.

